

UNOBSTRUCTEDNESS OF FILLING SECANTS AND THE GRUSON-PESKINE GENERAL PROJECTION THEOREM

ZIV RAN

ABSTRACT. The locus of multisequant rational curves to a variety of is nice when it fills up something nice.

1. INTRODUCTION AND STATEMENT OF THEOREM

Let P be a nonsingular, quasi-projective variety and $L \subset P$ a curve isomorphic to $\mathbb{P}^1$. Deformations of L as smooth curve in P are parametrized by some open subset G of a suitable Hilbert scheme. Let X be a nonsingular, locally closed subvariety of P . For a partition $(k.) = (k_1, \dots, k_r)$, L is said to be $(k.)$ -secant to X if the schematic intersection $L \cap X$ is a divisor of type $\sum k_i p_i$ on L , and *proper* as such if the p_i are pairwise distinct. The locus $S_{(k.)}$ of $(k.)$ -secant rational curves to X has a well-defined structure of locally closed subscheme of G , which may well be obstructed: i.e. singular, nonreduced or of dimension larger than the expected. It is therefore of interest to find conditions under which secant rational curves are unobstructed. Our purpose here is to prove such a condition. First a definition

Definition. Let Y be a smooth local germ along a smooth curve L . A local deformation $\mathcal{L} \rightarrow \text{Spec}(A)$ of L is said to be a *filling family* for Y if there exists an irreducible closed subscheme S of $\text{Spec}(A)$, such that the induced map $(\mathcal{L}_S)_{\text{red}} \rightarrow Y$ is dominant.

Theorem. *Let L be a complete nonsingular rational curve in P that is a proper $(k.)$ -secant to X and which moves in a family of $(k.)$ -secants filling up a smooth m -dimensional germ Y containing the germ of X at $L \cap X$. Then L is unobstructed as $(k.)$ -secant to X contained in Y . In particular, the scheme $S_{(k.),Y}$ of $(k.)$ -secants to X contained in Y is smooth at L of dimension equal to the expected, namely*

$$m - 3 - K_Y.L - kc + r, k := \sum k_i, c := m - n.$$

Moreover, for any partition $(l.)$ refining $(k.)$, the singularities of $S_{(l.),Y}$ at L are the same, up to a smooth factor, as those of the locus of divisors of type $(l.)$ on L at the divisor $L \cap X$.

Date: February 5, 2013.

1991 Mathematics Subject Classification. 14N05.

Key words and phrases. multisequant lines, rational curves.

The case where L is a line in $P = \mathbb{P}^N$ and Y is the germ of the ambient $\mathbb{P}^N$ itself is equivalent to the statement that the general projection of X to a lower projective space has the expected singularities, and was recently proven by Gruson and Peskine [2], settling a long-term conjecture, and extending many earlier partial results including [3] and [1]. This note grew out of an attempt to understand their exciting paper. Though our proof is independent of [2], our guiding principle is the following idea gleaned from it:

Uniformity principle: *The family of secants to X behaves in the same way at all points off X*

This principle is quite consequential: it implies, e.g. that if k -secants fill up the ambient space, then they are in general not $(k + 1)$ -secants, a statement nearly as strong as the theorem in the case $Y = \mathbb{P}^N$. Our idea is that this uniformity principle can be deduced from another, elementary and familiar, uniformity principle: namely that on a line, a generically spanned bundle is spanned; then, such a bundle has no higher cohomology either- just what we need for unobstructedness. In our case, the bundle in question is just the subsheaf of the normal bundle of a secant curve L as above, defined by the conditions of meeting X and being contained in Y . It is quite easy to check that all the required conditions are satisfied, which yields our result.

We thank Yeongrak Kim for helpful discussions. Hopefully, our indebtedness to the breakthrough work of Gruson and Peskine would be obvious to anyone.

2. FINITE SCHEMES ON A LINE

Let L be a nonsingular rational curve. Let $L^{(k)}$ denote the k -th symmetric product, or equivalently the degree- k Hilbert scheme of L . Then $L^{(k)}$ is canonically stratified by subschemes $L^{(k.)}$ for all partitions $(k.)$ of weight k , where $L^{(k.)} \subset L^{(l.)}$ iff $(l.)$ is a refinement of $k.$, given by a decomposition $k_i = \sum_j l_{i,j}$, and moreover $L^{(l.)}$ is reduced and is singular only along some proper substrata. Locally at an interior point $z = \sum k_i p_i \in L^{(k.)}$, we can decompose

$$(1) \quad L^{(k)} = \prod (\mathbb{A}^{k_i-1}) \times A$$

where each $\mathbb{A}^{k_i-1}$ is identified with the versal deformation of $k_i p_i$ and A is smooth, and $L^{(l.)}$ decomposes as $\prod D_i(l.) \times A$ where each $D_i(l.)$ is the subscheme of $\mathbb{A}^{k_i-1}$ corresponding to schemes of type $\sum_j l_{i,j} p_{i,j}$. In particular, when $(l.) = (k.)$, each D_i is a point.

All of this is well known (e.g. [4]) and follows from exact sequences

$$0 \rightarrow T_{L/z} \rightarrow T_L \rightarrow N'_{z,L} \rightarrow 0$$

$$0 \rightarrow N'_{z,L} \rightarrow N_{z,L} \xrightarrow{2} T_z^1 \rightarrow 0$$

where N denotes the lci normal bundle and N' is its 'locally trivial deformations' subsheaf (which is isomorphic to $J_Z/\mathcal{I}_Z$, J = Jacobian ideal, and in this case has length 1 at each p_i).

Let $S_{k,Y} = S_{(1^k),Y}$ where $(1^k) = (1, \dots, 1)$ of weight k . Then there is a locally defined classifying map

$$(2) \quad \mu : S_{k,Y} \rightarrow L^{(k)}$$

such that $S_{(k),Y} = \mu^{-1}(L^{(k)})$. Hence there is a map

$$(3) \quad \bar{\mu} : S_{k,Y} \rightarrow \prod (\mathbb{A}^{k_i-1})$$

whose special fibre is $S_{(k),Y}$ and such that

$$\bar{\mu}^{-1}(\prod D_i(l)) = S_{(l),Y}$$

3. THE SECANT AND CONTACT SHEAVES

Let notations be as in the Theorem. Then the normal bundle $N_{L/Y} \subset N_{L/P}$ is a semi-positive vector bundle on L , hence is a direct sum of line bundles of nonnegative degree, and therefore $H^1(N_{L/Y}) = 0$. It follows that the deformations of L in Y form a germ of a nonsingular subvariety of the Hilbert scheme parametrizing deformations of L in P , whose dimension equals $h^0(N_{L/Y}) = m - 3 - K_Y.L$.

Next, we make a local study of the subsheaf of the normal bundle corresponding to deformations of L preserving either the total length, or the exact type, of the intersection with X . Let $p \in X \cap L$ be a point of multiplicity k . We assume $k > 1$ as the case $k = 1$ is simpler. Choose local coordinates $x_1, \dots, x_N$ on P at p so that L is defined by $x_2, \dots, x_N$. We may choose local equations $g_1, \dots, g_c, c = m - n$, cutting out X on Y , and rearrange the x_i , so that

$$(4) \quad \begin{aligned} g_1 &\equiv x_1^k + x_N, \\ g_i &\equiv x_i, i = 2, \dots, c. \end{aligned}$$

modulo higher order terms. Then a local deformation of L in Y corresponds to setting

$$(5) \quad x_j = b_j(x_1), j \geq 2.$$

The condition that this deformation moves $X \cap L$ flatly is

$$b_j \equiv b_N \pmod{x_1^k}, 2 \leq j \leq c.$$

The further condition that the deformation also preserve the contact order near p is

$$(6) \quad b_N \equiv 0 \pmod{x_1^{k-1}}.$$

Applying this at each of the r intersection points $p_1, \dots, p_r \in X \cap L$, we conclude that there are subsheaves

$$(7) \quad N^c \subset N^s \subset N_{L/Y},$$

called the *contact* and *secant* sheaves, that parametrize motions of L within Y for which $L \cap X$ deforms flatly (for N^s) or locally trivially (for N^c), such that

$$(8) \quad \begin{aligned} \ell_{p_i}(N_{L/Y}/N^s) &= k_i(c-1), \\ \ell_{p_i}(N^s/N^c) &= k_i-1. \end{aligned}$$

Canonically, we have

$$(9) \quad \begin{aligned} N^s &= \text{Hom}(\mathcal{I}_L, \mathcal{O}_L) \cap \text{Hom}(\mathcal{I}_X \cap \mathcal{I}_L, \mathcal{I}_X \cdot \mathcal{O}_L), \\ N^c &= N^s \cap \text{Hom}(\mathcal{I}_L, J_{X \cap L, L}) \end{aligned}$$

where $\mathcal{I}$ denotes ideal in Y and J denotes Jacobian ideal (which parametrizes equisingular deformations). Note that N^s and N^c are respectively the sheaves classifying deformations of L preserving the length (resp. the local multiplicities) of $L \cap X$.

4. PROOF OF THEOREM

Our filling hypothesis means that N^c is generically spanned (a fortiori, so is N^s). Hence they are semipositive and have $H^1(N^s) = H^1(N^c) = 0$. The vanishing immediately implies the smoothness of $S_{k,Y}$ and of the local map $\bar{\mu}$ in (2), proving the Theorem.

REFERENCES

1. R. Beheshti and D. Eisenbud, *Fibers of generic projections*, Comp. math **146** (2010), 435–456.
2. L. Gruson and C. Peskine, *On the smooth locus of aligned hilbert schemes. The k -secant lemma and the general projection theorem*, Duke math. J. (2013), arxiv.org 1010.2399.
3. Z. Ran, *The (dimension+2)-secant lemma*, Invent. math **106** (1991), 65–71.
4. E. Sernesi, *Deformations of algebraic schemes*, Springer International, 2006.

ZIV RAN
 UNIVERSITY OF CALIFORNIA MATHEMATICS DEPARTMENT
 BIG SPRINGS SURGE FACILITY
 RIVERSIDE CA 92521
 ZIV.RAN@UCR.EDU